10 16525 Rec'd PCT/PTO 12 0CT 2000

				• • • • • • • • • • • • • • • • • • • •	ASSURE OF LOWLIN THOSE SOF			
FORM I	PTO-139 1-98)	Cr(Modified) U.S. DEPARTMI	ENT OF COMMERCE PATENT AND	FRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER			
TRANSMITTAL LETTER TO THE UNITED S			R TO THE UNITED	STATES	3027.00014			
		DESIGNATED/ELEC	TED OFFICE (DO/E	O/US)	U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR			
		CONCERNING A FIL	•	•	09/673558			
INTE		IONAL APPLICATION NO.	INTERNATIONAL FILI		PRIORITY DATE CLAIMED			
		PCT/CA99/00364	April 16,		April 16, 1998			
		NVENTION		6	IPE			
COM	IB-L	IKE SOLID PHASE FOR	MEASURING ANALY	TES /				
				OCT	1.2 man			
		r(s) for do/eo/us		2				
		. Marquardt		VEN.				
Znib	o Ga	n			MRADEMARIE.			
Appl	icant h	erewith submits to the United	States Designated/Elected O	ffice (DO/EO/US) th	e following items and other information:			
1.	\boxtimes	This is a FIRST submission of	of items concerning a filing	under 35 U.S.C. 371.				
2.		This is a SECOND or SUBS !	EQUENT submission of ite	ms concerning a filin	g under 35 U.S.C. 371.			
3.	\boxtimes	This is an express request to be	pegin national examination p	procedures (35 U.S.C	. 371(f)) at any time rather than delay 71(b) and PCT Articles 22 and 39(1).			
1	×	•			19th month from the earliest claimed priority date.			
4. 5.	×				19th month from the earnest claimed priority date.			
J.		A copy of the International Application as filed (35 U.S.C. 371 (c) (2))						
		 a. \(\subseteq \) is transmitted herewith (required only if not transmitted by the International Bureau). b. \(\subseteq \) has been transmitted by the International Bureau. 						
n		c. is not required, as the application was filed in the United States Receiving Office (RO/US).						
7		A translation of the Internation	• •		· ,			
Ü	×	A copy of the International Se	· · · · · · · · · · · · · · · · · · ·					
	×	Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))						
UT **	_	a. are transmitted herewith (required only if not transmitted by the International Bureau).						
		b. \(\square\) have been transmitted by the International Bureau.						
in de la companya de			; however, the time limit for		ments has NOT expired.			
, Kan			and will not be made.		•			
_9.		A translation of the amendme	ents to the claims under PCT	Article 19 (35 U.S.C	C. 371(c)(3)).			
10.	\boxtimes	An oath or declaration of the	inventor(s) (35 U.S.C. 371 (c)(4)).				
#1. #2.	\boxtimes	A copy of the International P	reliminary Examination Rep	ort (PCT/IPEA/409).				
12.		A translation of the annexes t	o the International Prelimina	ary Examination Rep	ort under PCT Article 36			
		(35 U.S.C. 371 (c)(5)).						
It		3 to 20 below concern docum						
13.	⊠	An Information Disclosure S						
14.		-		sheet in compliance	with 37 CFR 3.28 and 3.31 is included.			
15.	\boxtimes	A FIRST preliminary amend						
16.		A SECOND or SUBSEQUE	NT preliminary amendment		E OF MAILING BY "EXPRESS MAIL"			
17.		A substitute specification.	41 41 4	_	FL40557399643			
18.		A change of power of attorne		"EXPRESS MA	IL" Mailing Label Number			
19. 20	×	Certificate of Mailing by Express Mail Date of Deposit 10-12-00 Other items or information: I hereby certify that this capet or see is being income as being						
20.		Other items or information:		- with the limites	4 CONFE DOMEST CONTROL FOR STANDED TO SELECTION OF SELE			
1				Rock Office To	Addressee' scrvice under 07 CFR 1.10 dicated upping and is addressed to the			
ł				Assistant Com	missioner for Patents, Washington, D.C.			
}				20231.	2.1			
}				for	wh D			
		Marie M. Deut H						
		(Signature of person maling paner or fee)						
j								

532 Rec'd PCT/PTC 12 OCT 2000

J _{&} S. APPLICATION NO	O. (IF KNOWN SEE 27 CER 8	INTERNATIONAL APP PCT/CA99			1	00014
21. The follo	wing fees are submitted:.				CALCULATIONS	PTO USE ONLY
	FEE (37 CFR 1.492 (a) (1) -	(5)):			<u></u>	
☐ Neither international s and Internation	ational preliminary examination earch fee (37 CFR 1.445(a)(2) p nal Search Report not prepared	fee (37 CFR 1.482) nor paid to USPTO by the EPO or JPO	\$1,	000.00		
USPTO but In	oreliminary examination fee (37 sternation Search Report prepare	ed by the EPO or JPO	\$	860.00		
but internation	oreliminary examination fee (37 al search fee (37 CFR 1.445(a)	(2)) paid to USPIO		710.00		
but all claims	oreliminary examination fee pai did not satisfy provisions of PC	T Article $33(1)$ - (4)		690.00		
☐ International pand all claims	oreliminary examination fee pai satisfied provisions of PCT Ar	ticle 33(1)-(4)	. 1	s100.00 	1	
	ENTER APPROPRI	ATE BASIC FEE	AMOUNT	=	\$860.00	
Surcharge of \$130.00 nonths from the earli	for furnishing the oath or declariest claimed priority date (37 C	aration later than FR 1.492 (e)).	□ 20 □	30	\$0.00	
CLAIMS	NUMBER FILED	NUMBER EXTR				
Fotal claims	21 - 20 =	1	x \$1	8.00	\$18.00	
ndependent claims	3 - 3=	0	x \$8	0.00	\$0.00	
Multiple Dependent	Claims (check if applicable).				\$0.00	
		F ABOVE CALCU		=	\$878.00	
Reduction of 1/2 for must also be filed (N	filing by small entity, if applications of the small entity is application of the small entity of the small entities of the small entity of the small entity of the small entities of the small entity of the small entity of the small entities of the small entity of the small	able. Verified Small Entineck if applicable).	ty Statement	×	\$439.00	
			SUBTOTA	L =	\$439.00	
Processing fee of \$13 months from the earl	30.00 for furnishing the English iest claimed priority date (37 C	translation later than CFR 1.492 (f)).	□ 20 □	30 +	\$0.00	
		TOTAL NATION	ONAL FEE	=	\$439.00	
Fee for recording the	e enclosed assignment (37 CFR appropriate cover sheet (37 CFR	1.21(h)). The assignment 3.28, 3.31) (check if ap	t must be oplicable).		\$0.00	
- Erreit		TOTAL FEES I	ENCLOSEI		\$439.00	
					Amount to be: refunded	\$
17000 1 10000 10000 10000					charged	\$
A check in the amount of \$439.00 to cover the above fees is enclosed. Please charge my Deposit Account No. in the amount of to cover the above fees. A duplicate copy of this sheet is enclosed.						
to Deposit A	issioner is hereby authorized to Account No. 11-1449	A duplicate copy of this	s sheet is enclose	ed.		yn.
NOTE: Where an 1.137(a) or (b)) mu	appropriate time limit under st be filed and granted to rest	57 CFK 1.494 or 1.495 l ore the application to po	nas not been me ending status.	ei, a petitio	on to revive (3/ Cr	~ ()
SEND ALL CORRI	ESPONDENCE TO:				()	
Amy E. Rinaldo KOHN & ASSOC	IATES		SIGN	IATURE		
30500 Northweste	rn Hwy.		Amy	E. Rinal	do	···
Suite 410 Farmington Hills,	Michigan 48334		NAM	IE		
(248) 539-5050			45,7	91		
					N NUMBER	
				ber 12, 2		
			DAT			
			DAI	L		

. Attorney's Docket Number: _3	027.00014 PATENT
Applicant or Patentee:	Ronald R. Marquardt et al
Serial or Patent No:	
Filed or Issued:	
For:	COMB-LIKE SOLID PHASE FOR MEASURING
	ANALYTES
(37 CFR 1.9(f) and I hereby declare that I am an	CLARATION) CLAIMING SMALL ENTITY STATUS d 1.27(d) NONPROFIT ORGANIZATION official empowered to act on behalf of the
nonprofit organization identi	fied below:
Name of Organization: Mani	toba Cancer Treatment & Research Foundation
Organization Address: 100	Olivia, Room 0N206
Winn	ipeg, Canada R3E-OV9
Type of Organization:	
University or oth	er Institution of Higher Education
X Tax exempt under (26 USC 501(a) an	Internal Revenue Service Code d 501(c)(3)
	fic or Educational under Statute of, USA, Statute No
Service Code (26	Tax Exempt under Internal Revenue USC 501(a) and 501(c)(3) if located tes of America (State:) ute:)
qualifies as a nonprofit organ purposes of paying reduced fe	profit organization identified above nization as defined in 37 CFR 1.9(e) for es under Section 41(a) and 41(b) of Title h regard to the invention as entitled above.
Described in:	
X the specification	n filed herewith with title as listed above
the application :	identified above.

I hereby declare that rights under contract or law have been conveyed to and remain with the nonprofit organization with regard to the above identified invention.

the patent identified above.

If the rights held by the nonprofit organization are not exclusive, each individual, concern or organization having rights to the invention is listed below* and no rights to the invention are held by any person, other than the inventor, who could not qualify as a small business concern under 37 CFR 1.9(d) or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or nonprofit organization under 37 CFR 1.9(e).

Separate verified statements are required from each named person, concern

or orga as smal	nization having rights 1 entities.(37 CFR 1.2	to the invention averring to their status
	rson, concern or or listed below:	ganization having any rights in the
	No such person, co	oncern, or organization exists.
X	Each such person,	concern or organization is listed below.
NAME:	University of Mani	toba
ADDRESS:	Administration Bui	lding, Room 202
	Winnipeg, Manitoba	, Canada R3T-2N2
Individua	al Small Busi	ness <u>X</u> Nonprofit Organization
NAME:		
ADDRESS:		
Individua	al Small Busi	ness Nonprofit Organization
notification to small ent earliest of	of any change in ity status prior the issue fee or a	file, in this application or patent, status resulting in loss of entitlement to paying, or at the time of paying, the any maintenance fee due after the date on is no longer appropriate. [37 CFR 1.28(b)]
true and that to be true; knowledge th punishable by 18 of the Un- jeopardize th	t all statements ma and further tha nat willful false y fine or imprisonm ited States Code, a ne validity of the	ements made herein of my own knowledge are ade on information and belief are believed to these statements were made with the statements and the like so made are ment, or both, under Section 1001 of Title and that such willful false statements may application, any patent issuing thereon, ified statement is directed.
Name of Perso	on Signing:	Dr. Brent Schacter
Title in Orga	nization:	President/CEO
Address of Pe	erson Signing:	100 Olivia, Room ON206
		Winnipeg, Manitoba, Canada
SIGNATURE:	Tout So	Date: 10-12-00

Attorney's I	Oocket Number:	3027.00014 PATENT
Applicant or	Patentee:	Ronald R. Marquardt, et al
Serial or Pat	ent No.:	
Filed or Issu	ed:	
Title:		COMB-LIKE SOLID PHASE FOR MEASURING ANALYTES
	VERIFIED STATEM (37 CFR 1.9(f) and	ENT CLAIMING SMALL ENTITY STATUS 1.27(d) — NONPROFIT ORGANIZATION
I hereby declaridentified bel	are that I am an official ow:	empowered to act on behalf of the nonprofit organization
Name of Org	anization:	The University of Manitoba
Organization	Address:	Administration Building - Room 202
		Winnipeg, Manitoba, CANADA R3T 2N2
Type of Orga	nization:	
<u>X</u>	University or other Ins	stitution of Higher Education
	Tax exempt under Into (26 USC 501(a) and 50	ernal Revenue Service Code 01(c)(3)
	America State:	Educational under Statute of State of the United States of
	Would Qualify as Tax Service Code (26 USC America (State:	Exempt under Internal Revenue 501(a) and 501(c)3) if located int he United States of
organization a	lare that the nonprofi as defined in 37 CFR 1.9	t organization identified above qualified as a nonprofit (e) for purposes of paying reduced fees under Section 41(a) s Code with regard to the invention as entitled above.
Described in	:	
<u>X</u>	the specification filed l	herewith with title as listed above.
	the application identifi	ied above.
	the patent identified a	bove.
1		

I hereby declare that rights under contract or law have been conveyed to and remain with the nonprofit organization with regard to the above identified invention.

If the rights held by the nonprofit organization are not exclusive, each individual, concern or organization having rights to the invention is listed below* and no rights to the invention are held by any person, other than the inventor, who could not qualify as a small business concern under 37 CFR 1.9(d) or nonprofit organization under 37 CFR 1.9(3).

Separate verified statements are required from each person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

Each such pe	rson, concern or orga	nization having any	rights int h	e invention is listed below:
	No such person, con	cern, or organizatio	n exists.	
	Each such person, c	oncern or organizat	ion is listed	below.
NAME:				
ADDRESS:				
	Individual	Small Business	Nonp	rofit Organization
NAME:				
ADDRESS:				
	Individual	Small Business	Nonp	rofit Organization
earliest of the	ss of entitlement to sm	nall entity status pr ntenance fee due at	ior to paving.	tion of any change in status , or at the time of paying, the on which status as a small
statements m statements we punishable by Code, and tha	ade on information a re made with the know fine or imprisonment.	and belief are belie wledge that willful f , or both, under Sec atements may jeopa	eved to be trained alse statement to a local training to the the value of the value	wledge are true and that all rue; and further that these nts and the like so made are Title 18 of the United States lidity of the application, any ent is directed.
Name of Perso	on Signing:	M.W. McAdam		
Title in Organ	nization:	Vice-President, A	<u>lministratior</u>	1
		Winnipeg, Manito	ba, Canada	R3T 2N2
Signature:	mwind	<u></u>	Date:	October 12, 2000
	(Small Entity	Nonrestt Ferm 7	'21 Dogg 2	· -£2)

(Small Entity-Nonprofit [Form 7-3] — Page 2 of 2)

532 Rec'd PCT/PTG 12 0CT 2000

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

RONALD R. MARQUARDT, ET AL

Serial No.: Unknown

Filed: Herewith

For: COMB-LIKE SOLID PHASE FOR MEASURING ANALYTES

Our File No.: 3027.00014

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents Washington, D.C. 20231

Sir:

Please preliminarily amend the above-captioned patent application prior to examination on the merits as follows:

IN THE SPECIFICATION:

Page 1, after the Title, please insert the following section:

-- CROSS REFERENCE TO RELATED APPLICATIONS

This patent application is a National Phase Concerning a Filing Under 35 U.S.C. 371, claiming the benefit of priority of PCT/CA99/00364, filed April 16, 1999, which claims the benefit of priority of Canadian Application Serial No. 2,228,821, filed April 16, 1998, both of which are incorporated herein by reference.—

REMARKS

The above amendments add no new matter and are merely to claim benefit of priority.

It is respectfully submitted that the application is now in condition for allowance, which allowance is respectfully requested.

Respectfully submitted,

KOHN & ASSOCIATES

Amy E. Rinaldo Registration No. 45,791 30500 Northwestern Highway Suite 410 Farmington Hills, MI 48334 (248) 539-5050

CERTIFICATE OF MAILING BY "EXPRESS MAIL"

Express Mail Mailing Label No.: EL 405595996US Date of Deposit: October 12, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office To Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to: Assistant Commissioner for Patents, Washington, DC 20231, BOX PATENT APPLICATION.

Marie M DeWitt

532 Rec'd PCT/PTC 1.2 OCT 2000

COMB-LIKE SOLID PHASE FOR MEASURING ANALYTES

FIELD OF THE INVENTION

The present invention relates generally to the field of biological and biochemical assays.

BACKGROUND OF THE INVENTION

It is of great importance in all fields and disciplines of the life sciences to utilize qualitative and quantitative analytical techniques for the detection, identification, and measurement of a wide variety of biologically important molecules. For example, these analytical techniques can be utilized in many different types of assays including those for enzymes, receptors, lectins and inhibitors.

As virtually all chemical reactions in living systems are catalyzed by enzymes, the assay of enzyme activity is one of the most frequently encountered procedures in biochemistry. Typically, most enzyme assays are used to estimate the amount or activity of an enzyme present in a cell, tissue, other preparation, or as an essential part of an enzyme purification protocol. The current assay methods have been developed based on the physical, chemical and immunological properties of the enzyme of interest and use detection means such as for example photometric, radiometric, high performance liquid chromatographic, and electrochemical assays. (Eisenthal, R. and Danson, M. J., in *Enzyme Assays: a practical approach*. (IRL Press: Oxford University, Oxford, (1993)). Although the above-described methods basically fulfill the requirements for routine analysis, there are the disadvantages of low sensitivity (Oppert et al, 1997, *BioTechniques* 23:70-72), multiple steps (Twining, S. S., 1984 *Anal Biochem* 143: 30-344; Pazhanisamy, S. et al, 1995, *Anal Biochem* 229: 48-53) and steps that are tedious and time-consuming (Fields, R.,. 1972, *Meth Enz* 25B: 464-468).

Similarly, immunoassays have been widely using in human clinical tests and therapeutics, agriculture, food, veterinary and environmental diagnostics (Deshpandes, S. in *Enzyme immunoassays from concept to product development.*, (Chapman & Hall: New York, 1996)). While for most purposes, immunoassays are effective (Cleaveland, J. S. et al, 1990, *Anal Biochem* 190: 249-253.), in some cases they are not suitable, for example, in the determination of enzyme activity. This is due to the fact that binding assays for antibody and antigen (in this instance, an enzyme)

only measure the concentration of an antigen (enzyme) and not its activity. Specifically, it is often important to know the catalytic activity of an enzyme and not just the concentration of the enzyme as a given amount of enzyme may have varying activity depending on reaction conditions, source of enzyme and the isolation protocol used. Also, antibodies tend to react only with structurally similar antigens. Therefore, it is often not possible to quantitate the amount of an enzyme from a given species using specific antibodies from a related species using immunoassays.

The pharmaceutical industry utilizes the methods mentioned above to screen compounds for discovering drugs. This process is slow due to the multiple steps required and the large amount of compounds needed to be tested. Typically, on a good day, a lab might test 100 to 1,000 compounds. However, in the race to commercialize, pharmaceutical manufacturers are facing great pressure to reduce the time required to discover new clinical drugs, cut assay costs, and screen more compounds and against more targets. Therefore, there is a very high demand to develop new methods to meet the requirements of High Throughput Screening (HTS). There has been described a method using quenched BODIPY dye-labeled casein as a substrate for determining the activities of protease, which is sensitive and amenable to automation (Jones, L. J. et al, 1997, Anal Biochem 251: 144-152). The degree of guenching of the fluorescent tag is crucial in this method. However, if there is not enough quenching due to poor conjugation or degradation of the fluorescencelabeled substrate under storage, etc. the assay will not be very useful. Also this procedure has relatively high background values which reduce its sensitivity. Another example of a potentially useful high throughput assay was made by Marquardt, et al. and described in PCT/WO97/43438. The method involves many steps, including coating the wells of a microplate, washing the wells, adding biologically active substance to the wells, washing the wells once more, adding the indicator enzyme to the wells, washing the wells again and adding a color development reagent. As a result, the assay cannot be readily used in assays requiring rapid analysis.

Clearly, a new assay method that is highly sensitivity and suitable for high throughput assays is needed. Specifically, the assay should be suitable for a wide variety of applications, easy to use and require only a few steps for obtaining

results.

SUMMARY OF THE INVENTION

It is an object of the invention to provide a new assay method that is highly sensitive and suitable for high throughput assays.

This invention provides a new method for the qualitative and quantitative analysis of bioactive substances. The assay method is based on a onestep procedure for separating the reactants from the products (resultants) after completion of the reaction followed by measurement of the amount of a labeled reactant or labeled product that has been left in the reaction vessel or on the pin. The device consists of two parts; one of which is a reaction vessel and other is a pin-like device (a probe) that can fit into the reaction vessel. The reaction begins by insertion of the pin-like device that is coated with a reactant into the reaction vessel containing the same or other reactants than those coated on the pin-like surface. The said pinlike surface is taken out from the reaction vessel to stop the reaction. The amount of the labeled products or labeled reactant remaining in the reaction vessel or on the probe can be determined according to the intensity of its label which can be fluorescent, luminescent and chromogenic molecules or radioactive tag, etc. The amount of the label in the reaction vessel is directly or reciprocally proportional to the activity or amount of the bioactive substance that is to be measured. The device can be used for the assay of enzymes, their inhibitors, antibodies and antigens, and receptors or lectins and their ligands.

According to a first aspect of the invention, there is provided a method for measuring the activity or concentration of a biomolecule comprising:

providing a reaction vessel containing a sample, said sample including a biomolecule having a biological activity;

providing a probe coated with a reactant, said reactant being capable of interacting with the biomolecule;

adding a known quantity of a compound with a detectable label to the sample;

inserting the probe into the reaction vessel such that the biomolecule and the detectable label contact the reactant and interact with the reactant such that

label is bound to either the reactant or the biomolecule;

removing the probe from the reaction vessel; and

measuring the quantity of detectable label in the reaction vessel and/or on the probe.

The probe may have a shape selected from the group consisting of: pin-like; cone-like; cuboid; cylindrical; star-shaped; and spire-shaped.

The detectable label may be selected from the group consisting of: colorimetric label; radioactive label; luminescent label; and fluorescent label.

The sample may be a biological sample.

The biological activity may be an enzyme; a receptor; a lectin or an antibody; or a specific ligand of an antibody, a receptor, a lectin or an enzyme.

The reactant may be bound to the probe.

The sample may include an inhibitor of the biological activity of the biomolecule.

The sample may include a competitor of the biological activity of the biomolecule.

The biomolecule may be selected from the group consisting of: an enzymatic product; an enzyme; a substrate; a receptor; an inhibitor; a ligand; a lectin; a lectin-binding ligand; a receptor-binding ligand; an antibody; an antigen; a receptor; and a lectin.

The compound may be selected from the group consisting of: an enzymatic product; an enzyme; a substrate; a receptor; an inhibitor; a ligand; a lectin; a lectin-binding ligand; a receptor-binding ligand; an antibody; an antigen; a receptor; and a lectin.

According to a second aspect of the invention, there is provided a method for measuring the activity or concentration of a biomolecule comprising:

providing a reaction vessel containing a sample, said sample including a biomolecule having a biological activity;

providing a probe coated with a reactant, said reactant being capable of interacting with the biomolecule, said reactant including a detectable label;

inserting the probe into the reaction vessel such that the reactant and

the detectable label contact the biomolecule and interact with the biomolecule such that label is released from the reactant;

removing the probe from the reaction vessel; and

measuring the quantity of detectable label in the reaction vessel and/or on the probe.

The probe may have a shape selected from the group consisting of: pin-like; cone-like; cuboid; cylindrical; star-shaped; and spire-shaped.

The detectable label may be selected from the group consisting of: colorimetric label; radioactive label; luminescent label; and fluorescent label.

The reactant may be bound to the probe.

The sample may be a biological sample.

The biological activity may be an enzymatic activity or a binding affinity.

The sample may include an inhibitor of the biological activity of the biomolecule.

The sample may include a competitor of the biological activity of the biomolecule.

The biomolecule may be selected from the group consisting of: an an enzymatic product; an enzyme; a substrate; a receptor; an inhibitor; a ligand; a lectin; a lectin-binding ligand; a receptor-binding ligand; an antibody; an antigen; a receptor; and a lectin.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned hereunder are incorporated herein by reference.

DEFINITIONS

"Ligand" as used herein refers to a bioactive molecule having specific binding affinity for another biomolecule.

Stun dam

the first

The state of the s

WO 99/54734 PCT/CA99/00364

6

"Receptor" as used herein refers to a bioactive molecule that has a specific binding affinity for another biomolecule, for example, a ligand.

"Bioactive molecule" or "biologically active substance" as used herein except where otherwise stated refers to a molecule or complex having a biological activity, for example, an enzymatic activity or binding affinity for another biomolecule.

"Probe" as used herein refers to a member arranged to be reversibly inserted into a reaction mixture, as discussed below.

The present invention relates to a method for the detection, identification and measurement of the amount or activity of biologically active molecules via a one-step separation of reactants from products using a device comprising a probe whose surface is coated with a biologically active substance and a reaction vessel containing the reaction mixture. The reaction mixture contains a bioactive molecule and may, for example, be a biological sample or a synthetic reaction mixture in which all of the components are known.

As discussed above, the assay consists of a probe and a reaction vessel. As discussed above, the reaction vessel contains a reaction mixture which includes a bioactive molecule, for example, an enzyme or ligand, as well as possibly other compounds, for example, inhibitors, antibodies or receptors, as described below. The probe has a surface and the surface of the probe is coated with a compound. It is of note that the compound selected for coating is either a substrate for or has binding affinity for the bioactive molecule, as described below. Furthermore, the compound may be labeled, for example, fluorescently, luminescently, colorimetrically or radioactively, or the reaction mixture may contain a labeled substrate for or a labeled biomolecule having binding affinity for the bioactive molecule. In the first instance, wherein the compound is labeled, the labeled group on the compound is selected such that the activity of the bioactive molecule causes the label to be cleaved or removed from the compound and released to the reaction mixture. In the second instance, the label on the substrate within the reaction mixture is selected such that the activity of the bioactive molecule will cause the label to be transferred from the labeled substrate to the compound coating the surface of the probe, as described below. Alternatively, the labeled biomolecule and the biomolecule in the reaction mixture may compete for binding to the compound on the surface of the probe, or the biomolecule in the reaction mixture may compete with the labelled compound in the reaction mixture and the compound on the surface of the probe, as described below.

7

In general, the assay is carried out as follows: the quantity of labeled compound coated on the surface of the probe or the quantity of labeled substrate in the reaction vessel may first be determined by means known in the art according to the type of label used. The probe is then lowered into the reaction vessel such that the coated surface of the probe contacts the reaction mixture. The bioactive molecules present in the reaction mixture then either release label from the compound, transfer label from the substrate in the reaction mixture to the compound on the surface of the probe or compete for binding, as described below. Once the probe has been in contact with the reaction mixture for a pre-determined period of time, the probe is removed from the reaction mixture, thereby terminating the reaction without the need to add additional chemicals, heat the reaction mixture or on the surface of the probe can then be determined, which can then be used to calculate reaction kinetics.

In the embodiment described herein, the probe has a substantially pinlike shape. It is of note that in other embodiments, the probe may be of other shapes, for example, cylindrical, spire-shaped, star-shaped, cuboid, or cone-shaped.

This is in contrast with traditional methods, which, for example, may involve providing a reaction vessel containing a reaction mixture including a bioactive molecule, adding a labeled compound that is a substrate for the bioactive molecule, stopping the reaction, that is, stopping the biological activity of the bioactive molecule, and separating unused substrate from used substrate by washing or chromatography. An alternative type of assay known in the art involves coating a reaction vessel with a compound, adding a ligand, allowing binding between the ligand and the compound to occur, washing away unbound ligand, and adding a reporter compound to detect bound ligand.

Clearly, these methods are less desirable than the above-described

assay, as they require multiple steps and include washing and/or separation steps which can cause considerable inaccuracy in experimental results. However, the above-described assay does not require washing or separation steps and involves relatively few steps, resulting in shorter assay times, more reliable results and greater ease of use.

The invention will now be described using examples. Specifically, the examples provide broad examples of methods of using the assay as well as some specific examples. However, the scope of the invention is not limited to these examples

EXAMPLE I – TRANSFER OF LABEL FROM THE REACTION MIXTURE TO THE PROBE

For example, the surface of the probe, in this embodiment a pin-like device, is coated with reactant 1 which is an acceptor of a moiety from reactant 3. Reactant 3 is a substrate for reactant 2 having a biological activity. That is, in this example, reactant 1 is the compound used to coat the probe, reactant 3 is the labeled substrate and reactant 2 is the bioactive molecule. Furthermore, reactant 2 and reactant 3 are present in the reaction mixture in the reaction vessel. When the coated pin-like device is inserted into a reaction vessel, the labeled moiety from reactant 3 is transferred to reactant 1 due to the biological activity of reactant 2 acting on reactants 1 and 3. The amount of labeled reactant 3 remaining in the reaction vessel can be directly determined without an additional step and is reciprocally proportional to the activity of reactant 2 after the pin-like device containing the labeled moiety from reactant 3 has been taken out of the reaction vessel. Alternatively, the amount of label transferred to the probe may be determined, which is directly proportional to the activity of reactant 2.

EXAMPLE II - TRANSFER OF LABEL FROM THE REACTION MIXTURE TO THE PROBE IN PRESENCE OF INHIBITOR

In this example, the transfer of the labeled moiety between reactant 1 and reactant 3 by reactant 2 is interfered with by reactant 4 which is an inhibitor of reactant 2. Thus, when the coated probe is inserted into the reaction vessel, reactant 4 inhibits transfer of the labeled moiety from reactant 3 to reactant 1. The amount of

labeled reactant 3 remaining in the reaction vessel once the probe has been removed from the reaction vessel is directly proportional to the amount of reactant 4 (inhibitor) in the reaction mixture. Similarly, the amount of labeled reactant 1 present on the surface of the probe is reciprocally proportional to the amount of reactant 4 (inhibitor) present in the reaction mixture.

EXAMPLE III – TRANSFER OF LABEL FROM THE PROBE TO THE REACTION MIXTURE

In this example, the surface the probe, in this embodiment, a pin-like device is coated with reactant 1 which is a labeled substrate for reactant 2 having a biological activity. That is, in this example, reactant 1 is the labeled compound and reactant 2 is the bioactive molecule. Insertion of the pin-like device containing reactant 1 into a reaction vessel containing reactant 2 and other compounds essential for the reaction results in the initiation of a reaction in which labeled products of the reaction are released into the reaction vessel. The amount of the released label in the reaction vessel can be directly measured after the pin-like device has been taken out of the reaction vessel without an additional step and the quantity of label released is proportional to the activity of reactant 2. Alternatively, as discussed above, the amount of label remaining on the pin-like device can be determined, which would be reciprocally proportional to the activity of reactant 2.

EXAMPLE IV - TRANSFER OF LABEL FROM THE PROBE TO THE REACTION MIXTURE IN THE PRESENCE OF AN INHIBITOR

In this example, the release of label from reactant 1 coated on the surface of the pin-like device by reactant 2 is interfered with by reactant 3 which is an inhibitor of reactant 2 and is present in the reaction mixture. Insertion of the pin-like device containing reactant 1 into a reaction vessel containing reactant 2 and other compounds essential for the reaction as well as reactant 3, the inhibitor, results in the initiation of a reaction in which labeled products of the reaction are released into the reaction vessel. The amount of the released label from reactant 1 to the reaction vessel is reciprocally proportional to the amount of reactant 3 (inhibitor) after the pin-like device has been taken out of the reaction vessel. Alternatively, as discussed above, the amount of label remaining on the pin-like device can be determined,

which would be directly proportional to the activity of reactant 3 (inhibitor) on reactant 2 (bioactive molecule).

EXAMPLE V - CAPTURE OF ENZYMATIC PRODUCT BY THE PROBE

In this example, the surface of a pin-like device is coated with reactant 1 which is a binding ligand for product 1. Product 1 is formed from reactant(s) 3 linked with either a label or a binding agent capable of binding specifically to reactant 1. That is, product 1 is caused by the biological activity of reactant 2 in a reaction vessel acting on reactant(s) 3. The coated pin-like device is then inserted into the reaction vessel containing product 1 and other non-reacted reactants. Thus, the probe is coated with a compound (reactant 1) that binds or has binding affinity for a second compound (product 1). This second compound (product 1) is formed through the action of the bioactive molecule (reactant 2) and components of the reaction mixture (reactant(s) 3). Thus, as product 1 is formed through the action of reactant 2. product 1 binds to reactant 1 which coats the surface of the probe. After the pin-like device has been taken out of the reaction vessel, the amount of labeled non-reacted reactant 3 remaining in the reaction vessel can be directly measured without an additional step and is reciprocally proportional to the activity of reactant 2. Alternatively, the amount of labeled product 1 bound to the probe can be directly measured, which is directly proportional to the activity of reactant 2.

EXAMPLE VI – CAPTURE OF ENZYMATIC PRODUCT BY THE PROBE IN THE PRESENCE OF INHIBITOR

In this example, the formation of product 1 from reactants 3 due to biological activity of reactant 2 is interfered with by reactant 4 which is an inhibitor of reactant 2. The assay is carried out as described above (Example V), except that in this instance, the amount of labeled non-reacted reactant 3 remaining in the reaction vessel after the pin-like device has been taken out of the reaction vessel is directly proportional to the amount of reactant 4 (inhibitor) present in the reaction mixture. Alternatively, the amount of labeled product 1 bound to the probe can be directly measured, which is reciprocally proportional to the amount of reactant 4 (inhibitor) present in the reaction mixture.

EXAMPLE VII - COMPETITIVE BINDING

In this example, the surface of a pin-like device is coated with reactant 1 which is a binding ligand for reactant 3 being the labeled form of reactant 2 or a labeled compound that will compete for binding with reactant 2 for binding to reactant 1, as described below. Alternatively, reactant 3 may be chosen such that reactant 1 and reactant 2 will compete for binding to reactant 3, as described below. Competition between reactant 2 and reactant 3 for binding to reactant 1 or between reactant 2 and reactant 1 for binding to reactant 3 are initiated when the pin-like device bound using reactant 1 is inserted into a reaction vessel containing a unknown amount of reactant 2 and a known amount of labeled reactant 3. That is, the probe is coated with a first compound and the reaction vessel contains an unknown quantity of a second compound. A known quantity of a labeled third compound is added to the reaction vessel and the coated probe is inserted into the reaction mixture. Specifically, the labeled third compound is selected such that the third compound competes with the second compound for binding to the first compound or the first compound and the second compound compete for binding of the labeled third compound. The amount of the labeled reactant 3 remaining in the reaction vessel can be directly measured without an additional step after the pin-like device has been taken out of the reaction vessel and is directly proportional to the amount of reactant 2 in the reaction vessel. Alternatively, the amount of labeled product 1 bound to the probe can be directly measured, which is reciprocally proportional to the amount of reactant 2 in the reaction vessel. Thus, in the first instance labeled reactant 3 competes with reactant 2 for binding to the coated probe. In the second instance, reactant 1 and reactant 2 compete for binding to the labeled third compound. However, in both instances, the relative amount of reactant 3 bound to the probe or remaining in the reaction vessel provides information on the amount of reactant 2 in the reaction vessel.

EXAMPLE VIII - COMPETITIVE BINDING IN PRESENCE OF INHIBITOR

In this example, the reactions of the competitive binding between reactant 2 and reactant 3 for reactant 1 or between reactant 2 and reactant 1 for reactant 3 as described above are interfered with by reactant 4 which is an inhibitor of reactant 2. The amount of the labeled reactant 3 remaining in the reaction vessel

after the pin-like device has been taken out of the reaction vessel is directly proportional to the amount of reactant 4 (inhibitor) in the reaction vessel.

EXAMPLE IX - PROTEIN KINASE A ASSAY

In this example, an assay of protein kinase A activity in the presence and absence of its inhibitor using the above-described assay is described. As will be seen, this is an example of the assay described in Examples I and II, that is, the transfer of label from the reaction mixture to the probe in the presence and absence of inhibitor.

Materials: hydrolyzed and partially dephosphorylated casein, protein kinase 3': 5'-cyclic AMP dependent (PKA), cyclic AMP, protein kinase inhibitor, Na₂HPO₄, NaH₂PO₄, NaCl, Tween-20 are from Sigma. ³²P-ATP is from New England Nuclear. The 96-well microplate and 96-pin lid are from VWR Canlab.

Method:

1. Preparation of a coated 96-pin lid:

- (1). 5 g skim milk powder dissolved in 100 ml phosphate buffer saline (PBS, pH 7.2) is added to the wells of a microplate and incubated at 37° C for 2 hr. The microplate is washed three times with PBST (PBS + Tween).
- (2). Casein is dissolved in PBS to the concentration 5 ug/ml and 100 ul/well is added to each well of the microplate blocked with skim milk. The 96-pin lid is inserted into the wells of the microplate and incubated at 37° C for 3 hr, and is then taken out and rinsed with PBST. As a result of this arrangement, the probe is coated with casein.

2. Protein kinase assay:

- (1). A series of concentrations of protein kinase in phosphate buffer (PB pH 7.2, cAMP, ³²P-ATP) are added to the wells of a microplate (100 ul/well).
- (2). The reaction is initiated by inserting the 96-pin lid coated with casein. During incubation at 37° C for 30 min, ³²P is transferred from ³²P-ATP to casein by protein kinase. The 96-pin lid is then taken out of the wells of the microplate to stop the reaction.
- (3). The radioactivity of ³²P-ATP remaining in the wells of the microplate is counted in a scintillation counter and is reciprocally proportional to the activity of the

protein kinase.

- 3. Protein kinase inhibitor assay:
- (1). Varying concentrations of an inhibitor in the phosphate buffer (PB pH 7.2, cAMP, ³²P-ATP) are added to the wells (50 ul/well) of a microplate. Negative and positive controls are also included.
- (2). A fixed amount of protein kinase A (50 ul/well) is added to each well containing inhibitor and the controls. Inhibition of protein kinase activity by the inhibitor will occur.
- (3). The reaction of phosphate transferring is initiated by inserting the 96-pin lid coated with casein followed by incubation at 37° C for 30 min. Protein kinase catalyzes the transfer of ³²P from ³²P-ATP to casein with the degree of transfer being reciprocally related to the concentration of the inhibitor. The 96-pin lid is taken out of wells of the microplate to end the reaction.
- (4). The radioactivity of ³²P-ATP remaining in the wells of the microplate is counted in a scintillation counter and is directly proportional to the amount of the inhibitor.

EXAMPLE X - FLUORESCENT ASSAY OF PROTEASE AND PROTEASE INHIBITORS

In this example, the activity of a variety of proteases and protease inhibitors are examined using the above-described assay. As will be seen, this is an example of the general assay given in Examples III and IV, wherein the label is transferred from the probe to the reaction mixture.

Materials: proteinase K, elastase, protease XIII, papain, trypsin, pepsin, casein, dimethyl sulfoxide (DMSO), Na₂HPO₄, NaH₂PO₄, NaCl, Tween-20, citrate, ovomucoid, aprotinin are from Sigma. NHS-coumarin is from Molecular Probes. The 96-well microplate and 96-pin lid are from VWR Canlab.

Method:

- 1. Preparation of a fluo-casein: 5 mg NHS-coumarin in 100 ul DMSO is mixed with 10 mg casein in PBS (pH 7.2) in a microcentrifuge tube and incubate at room temperature for 3 hr.
- 2. Preparation of a coated 96-pin lid:

- (1). 5 g skim milk powder dissolved in 100 ml PBS (pH 7.2) is added to the wells of a microplate and incubated at 37° C for 2 hr. The microplate is washed three times with PBST.
- (2). Fluo-casein is dissolved in PBS to the concentration 5 ug/ml and 100 ul/well is added to each well of the microplate blocked with skim milk. The 96-pin lid is inserted into the wells of the microplate, incubated at 37° C for 3 hr, then taken out and rinsed with PBST.

3. Protease activity assay:

- (1). A series of concentrations of a protease in a buffer (100 ul/well) are added to the wells of a microplate.
- (2). The reactions are initiated by inserting the 96-pin lid coated with the fluo-casein and incubated at 37° C or room temperature for 30 min. The fluo-casein is hydrolyzed by protease to release the fluorescent labels into the reaction vessel. The 96-pin lid is taken out of the reaction vessel to stop the reaction.
- (3). The fluorescent intensity of the label in the wells of the microplate is measured with a fluorometer and is directly proportional to the activity of the protease.

4. Protease inhibitor assay:

- (1). Varying amounts of the protease inhibitor in the buffer (50 ul/well) are added to the wells of a microplate. Negative and positive controls are included.
- (2). A fixed concentration of the protease in the buffer (50 ul/well) is added to the wells containing inhibitor and the controls. The inhibition of the protease by the inhibitor reduces the amount of label released into the reaction mixture.
- (3). Hydrolysis of the casein by the protease is initiated by inserting the 96-pin lid coated with fluo-casein. The sample is incubated at 37⁰ C or room temperature for 1 hr. The fluo-casein is cleaved by the residual activity of the protease after inhibition to release the fluorescent label into the reaction vessels. The 96-pin lid is taken out of the reaction vessel to stop the reaction.
- (4). The fluorescent intensity of the label in the wells of the microplate is measured with a fluorometer and is reciprocally proportional to the amount of the inhibitor.

EXAMPLE XI - TELOMERASE ASSAY

In this example, telomerase activity, in the presence and absence of inhibitor is assayed. As will be seen, this is an example of Examples V and VI, binding of a synthesized product to the probe. Specifically, telomerase produces a nucleic acid molecule using labeled nucleotides and a primer that includes a binding agent which binds to the compound coating the probe. Thus, in this example, unextended primer will bind to the probe but will not produce a signal and unincorporated nucleotides will not be able to bind the probe and will remain in the reaction mixture.

Materials: Tris-acetate buffer (pH 8.5), potassium acetate, β -mercaptoethanol, spermidine, MgCl₂, EDTA, streptavidin are from Sigma. dATP, dTTP, fluo-dGTP, telomerase S10, RNase,

biotin-oligonucleotide primer may be purchased from Boehringer Mannheim or elsewhere. Inhibitor (7-deaza-dATP).

Method:

1. Preparation of a coated 96-pin lid:

- (1). 5 g skim milk powder dissolved in 100 ml PBS (pH 7.2) is added to the wells of a microplate and incubated at 37° C for 2 hr. The microplate is washed three times with PBST.
- (2). Streptavidin or avidin is dissolved in a buffer to the concentration 5 ug/ml and 100 ul/well is added to each well of the microplate blocked with skim milk. The 96-pin lid is inserted into the wells of the microplate and incubated at 37° C for 3 hr, and is then taken out and rinsed with PBST.

2. Telomerase activity assay:

(1). A series of concentration of telomerase in a reaction mixture containing 50 mM Tris-acetate pH 8.5, 50 mM potassium acetate (KAc), 5 mM β -mercaptoethanol, 1mM spermidine, 1mM MgCl₂, 0.5-2 mM dATP, 0.5-2 mM dTTP, 1.5 uM fluo-dGTP, 1 uM biotin-oligonucleotide primer (TTAGG)₃, are added to wells of a microplate and the mixture is incubated at 30° C for 1 hr. The reaction of DNA synthesis is stopped by adding the stop solution (10 mM Tris-HCl, pH7.5, 230 mM EDTA and 100 ug/ml RNase) at 37° C for 15 min.

- (2). Isolation of the fluorescent label-DNA with biotin-primer (fluo-DNA-biotin) from the reaction vessel is done by the binding of the fluo-DNA-biotin complex to the streptavidin (avidin)-coated 96 pin lid after the 96-pin lid has been inserted into the reaction vessel.
- (3). The fluorescent intensity of fluo-dGTP remaining in the reaction vessel is measured with a fluorometer after the 96-pin lid has been taken out of the reaction vessel and is reciprocally proportional to the activity of telomerase.

3. Telomerase inhibitor assay:

- (1). Varying amounts of the inhibitor in the reaction mixture (50 ul/well) are added to the wells of a microplate.
- (2). A fixed activity of telomerase in the reaction mixture (50 ul/well) is added to the wells containing an inhibitor and the controls. Incubate the wells of the plate at 30°C for 1-2 hr and stop the DNA synthesis with the stop solution.
- (3). Isolation of the fluorescent label-DNA with biotin-primer (fluo-DNA-biotin) from the reaction vessel is done via the binding of fluo-DNA-biotin complex to the streptavidin (avidin)-coated 96 pin lid after the 96-pin lid has been inserted into the reaction vessel.
- (4). The fluorescent intensity of fluo-dGTP remaining in the reaction vessel is measured with a fluorometer after the 96-pin lid has been taken out of the reaction vessel and is directly proportional to the amount of the inhibitor.

EXAMPLE XII - COMPETITIVE BINDING ASSAY

In this example, competitive binding of E. coli K88 fimbriae, receptor and inhibitor is assayed using the above-described method. This assay is a specific example of the general assay given in Examples VII and VIII, that is, of competitive binding assays using the above-described assay method.

Materials: Na₂HPO₄, NaH₂PO₄, NaCl, dimethyl sulfoxide (DMSO) are from Sigma. NHS-fluorescein is from Molecular Probes. A mucus receptor from a piglet and E. coli K88 fimbriae are prepared in a laboratory. Inhibitors are from different sources.

Method:

1. Preparation of fluo-fimbriae: 5 mg NHS-fluorescein in 100 ul DMSO is mixed with

3 % L w

10 mg fimbriae in 1 ml PBS (pH 7.2) followed by incubation at room temperature for 3 hr.

2. Preparation of a coated 96-pin lid

- (1). 5 g skim milk powder dissolved in 100 ml PBS (pH 7.2) is added to the wells of a microplate and incubated at 37° C for 2 hr. The microplate is washed three times with PBST.
- (2). Receptor is dissolved in PBS (pH 7.2) to a concentration of 5 ug/ml and 100 ul/well is added to each well of the microplate blocked with skim milk. The 96-pin lid is inserted into the wells of the microplate and incubated at 37⁰ C for 3 hr, and is then taken out and rinsed with PBST.

3. Receptor assay:

- (1). A series of concentrations of the receptor in a buffer (50 ul/well) are added to the wells of a microplate.
- (2). A fixed amount of the fluo-fimbriae (50 ul/well) is added to each well containing the receptor and the control.
- (3). Competitive binding reactions between the immobilized receptor and the free receptor (competitor) to the fluo-fimbriae are initiated by inserting the 96-pin lid coated with the receptor into the wells. Incubate at 37° C for 1 hr . Then the 96-pin lid is taken out of the vessel to stop the reaction.
- (4). The fluorescent intensity of the fluo-fimbriae remaining in the wells of the microplate is determined using a fluorometer and is directly proportional to the amount of the receptor (competitor).

4. Fimbriae or E. coli numeration assay:

- (1). A series of concentration of the fimbriae or E. coli cell in a buffer (50 ul/well) are added to the wells of a microplate.
- (2). A fixed amount of the fluo-fimbriae (50 ul/well) is added to each well containing the fimbriae or E. coli and control wells.
- (3). Competitive binding reactions between the fimbriae or E. coli (competitor) and the fluo-fimbriae to the immobilized receptor are initiated by inserting the 96-pin lid coated with the receptor into the wells. Incubate at 37⁰ C for 1 hr. Then the 96-pin lid is taken out of the vessel to stop the reaction.

1 h 1 r

- (4). The fluorescent intensity of the fluo-fimbriae remaining in the wells of the microplate is determined using a fluorometer and is directly proportional to the amount of the fimbriae or E. coli cells (competitor).
- 5. Inhibitor of the fimbriae-receptor binding assay:
- (1). A series of concentration of the inhibitor in a buffer (50 ul/well) are added to the wells of a microplate.
- (2). A fixed amount of the fluo-fimbriae (50 ul/well) is added to each well containing the inhibitor and the controls.
- (3). The reactions between the inhibitor and the immobilized receptor for binding to fluo-fimbriae or between the inhibitor and the fluo-fimbriae for binding to the immobilized receptor are initiated by inserting the 96-pin lid coated with the receptor into the wells. After incubation at 37° C for 1 hr the 96-pin lid is taken out of the vessel to stop the reaction.
- (4). The fluorescent intensity of the fluo-fimbriae remaining in the wells of the microplate is determined using a fluorometer and is directly proportional to the amount of the inhibitor (competitor).

As can be seen, the above-described assay has a wide range of applications due to its versatility. Furthermore, the assay does not require washing or "stopping" steps which are time-consuming and can lead to loss of signal or contamination. It is of note that in the examples above, the probe comprises a pin-shaped member and the reaction vessels are wells of a microtiter plate. However, it is to be understood that the probe may be of suitable shapes and arrangements, as discussed above, as may the reaction vessels.

While the preferred embodiments of the invention have been described above, it will be recognized and understood that various modifications may be made therein, and the appended claims are intended to cover all such modifications which may fall within the spirit and scope of the invention.

CLAIMS

1. A method for measuring the activity or concentration of a biomolecule comprising:

providing a reaction vessel containing a sample, said sample including a biomolecule having a biological activity;

providing a probe coated with a reactant, said reactant being capable of interacting with the biomolecule;

adding a known quantity of a compound with a detectable label to the sample;

inserting the probe into the reaction vessel such that the biomolecule and the detectable label contact the reactant and interact with the reactant such that label is bound to the reactant;

removing the probe from the reaction vessel; and

measuring the quantity of detectable label in the reaction vessel and/or on the probe.

- The method according to claim 1 wherein the probe has a shape selected from the group consisting of: pin-like; cone-like; cuboid; cylindrical; star-shaped; and spire-shaped.
- 3. The method according to claim 1 wherein the detectable label is selected from the group consisting of: colorimetric label; radioactive label; luminescent label and fluorescent label.
- 4. The method according to claim 1 wherein the reactant is bound to the probe.
- 5. The method according to claim 1 wherein the sample is a biological sample.
- 6. The method according to claim 1 wherein the biological activity is an enzymatic activity.
- 7. The method according to claim 1 wherein the biological activity is a binding affinity.
- 8. The method according to claim 1 wherein the sample includes an inhibitor of the biological activity of the biomolecule.

- 9. The method according to claim 1 wherein the sample includes a competitor of the biological activity of the biomolecule.
- 10. The method according to claim 1 wherein the biomolecule is selected from the group consisting of: an enzymatic product; an enzyme; a substrate; a lectin; a lectin-binding ligand; a receptor; an inhibitor; a receptor binding ligand; an antigen; and an antibody.
- 11. The method according to claim 1 wherein the compound is selected from the group consisting of: an enzymatic product; an enzyme; a lectin; a lectin-binding ligand; a substrate; a receptor; an inhibitor; a receptor binding ligand; an antigen; and an antibody.
- 12. A method for measuring the activity or concentration of a biomolecule comprising:

providing a reaction vessel containing a sample, said sample including a biomolecule having a biological activity;

providing a probe coated with a reactant, said reactant being capable of interacting with the biomolecule, said reactant including a detectable label;

inserting the probe into the reaction vessel such that the reactant and the detectable label contact the biomolecule and interact with the biomolecule such that label is released from the reactant:

removing the probe from the reaction vessel; and

measuring the quantity of detectable label in the reaction vessel and/or on the probe.

- 13. The method according to claim 12 wherein the probe has a shape selected from the group consisting of: pin-like; cone-like; cuboid; cylindrical; star-shaped; and spire-shaped.
- 14. The method according to claim 12 wherein the detectable label is selected from the group consisting of: colorimetric label; radioactive label; luminescent_label; and fluorescent label.
- 15. The method according to claim 12 wherein the reactant is bound to the probe.
 - 16. The method according to claim 12 wherein the sample is a

biological sample.

- 17. The method according to claim 12 wherein the biological activity is an enzymatic activity.
- 18. The method according to claim 12 wherein the biological activity is a binding affinity.
- 19. The method according to claim 12 wherein the sample includes an inhibitor of the biological activity of the biomolecule.
- 20. The method according to claim 12 wherein the sample includes a competitor of the biological activity of the biomolecule.
- 21. The method according to claim 12 wherein the biomolecule is selected from the group consisting of: an enzymatic product; an enzyme; a substrate; a receptor; a receptor ligand; an antigen; a lectin; a lectin-binding ligand; a ligand; and an antibody.

Docket No. 3027.00014

Declaration and Power of Attorney For Patent Application

English Language Declaration

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

COMB-LIKE SOLID PHASE FOR MEASURING ANALYTES

th	ne specification of which			
	check one)			
	is attached hereto.			
	was filed on	as Ur	nited States Application No.	or PCT International
134	Application Number			
	and was amended on _			
TT a			(if applicable)	
la I Tu ir	hereby state that I have rencluding the claims, as ame		the contents of the above in referred to above.	dentified specification,
	Office all information Federal Regulations,			
S a li: ir	section 365(b) of any foreing PCT International applies ted below and have also	ign application(s) for pat cation which designated identified below, by chec International application	e 35, United States Code, tent or inventor's certificate at least one country other the sking the box, any foreign ap a having a filing date before	or Section 365(a) of nan the United States, oplication for patent or
P	Prior Foreign Application(s)			Priority Not Claimed
2,2	228,821	Canada	16 April 1998	
	Number) CT/CA99/00364	(Country) Canada	(Day/Month/Year Filed) 16 April 1999	
(1	Number)	(Country)	(Day/Month/Year Filed)	
(I	Number)	(Country)	(Day/Month/Year Filed)	

I hereby claim the benefit unde application(s) listed below:	r 35 U.S.C. Section 119(e)	of any United States provisional
(Application Serial No.)	(Filing Date)	
(Application Serial No.)	(Filing Date)	
(Application Serial No.)	(Filing Date)	
Section 365(c) of any PCT Internations insofar as the subject matter of earlies United States or PCT International U.S.C. Section 112, I acknowledge Office all information known to make Section 1.56 which became available or PCT International filing date of the	tional application designating ach of the claims of this app application in the manner presented to the Leto be material to patentabile between the filing date of the contents of the contents.	any United States application(s), or the United States, listed below and, lication is not disclosed in the prior rovided by the first paragraph of 35 United States Patent and Trademark lity as defined in Title 37, C. F. R., the prior application and the national
(Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned)
(Application Serial No.) (Application Serial No.) (Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned)
(Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Cenneth I. Kohn (30,955)	ist name and registration number)	moldo (45 701)
tennem 1. Ronn (50,955)	, Amy E. Ri	naldo (45,791)
	Amy E Dinalda	
end Correspondence t		
	Kohn & Associates 30500 Northwestern Hwy. Suite 410	
	Farmington Hills, Michigan 48334	
	o: (name and telephone number)	
my E. Rinaldo (248) 539-5	050	
1		
Full name of sole or first inven		
Ronald Marquardt.	(C)	
Sole onfirst inventor's signatur	• 44 0 11	Date
K. K-Marge	all (A)	October <u>26</u> , 2000
Residence V 878 Kilkenny Drive Win	nipeg, Manitoba R3G 4G3 CANADA	
Citizenship	npeg, Maintoba K3G 4G3 CANADA	
CA		
Post Office Address		
Same as above.		
1		
210		
Full name of second inventor, Zhibo Gan	f any	
Second Inventor's signature		
Second inventor's signature	\sim \sim \sim	Date
Residence		Nov 1, Loon
212-99 Dalhousie Drive, V	Vinnipeg, Manitoba R3G 3M2 CANADA	
Citizenship	11331	
CA Post Office Address		